

## YM AGAR (7525)

### Intended Use

YM Agar is used for the cultivation of yeasts, molds, and other aciduric organisms.

### Product Summary and Explanation

YM Agar is prepared according to the formulation published by Wickerham.<sup>1-3</sup> YM Agar was formulated for the selective isolation of yeasts from mixed cultures containing bacteria and molds. Media selectivity may be enhanced through acidification or through addition of selective agents. YM Agar should be sterilized without pH adjustment and sterile acid added to the medium cooled to 45 - 50°C. Acidified YM Agar should not be heated. Antibiotics may be aseptically added to sterile media. Other fungistatic materials may be added to YM Agar to eliminate molds and permit enumeration of yeasts to mixed populations.

### Principles of the Procedure

Enzymatic Digest of Gelatin is a nitrogen and amino acid source in YM Agar. Yeast Extract provides trace elements and vitamins. Malt Extract is a source of carbon, protein, and nutrients. Dextrose is an energy source. Agar is the solidifying agent.

### Formula / Liter

Enzymatic Digest of Gelatin .....	5 g
Yeast Extract .....	3 g
Malt Extract .....	3 g
Dextrose .....	10 g
Agar .....	20 g

Final pH: 6.2 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

### Precaution

1. For Laboratory Use.

### Directions

1. Suspend 41 g of the medium in one liter of purified water.
2. Heat with frequent agitation and boil for one minute to completely dissolve the medium.
3. Autoclave at 121°C for 15 minutes.

### Optional

If desired, acidify YM Agar to pH 3.0 – 4.0 by adding sterile 10% HCl, Tartaric Acid, or 10% Citric Acid. Selective agents, e.g., penicillin (20 units per mL final concentration) or streptomycin (40 micrograms per mL final concentration) may be added to the medium after sterilization using aseptic technique.

### Quality Control Specifications

**Dehydrated Appearance:** Powder is homogeneous, free-flowing, and light beige.

**Prepared Appearance:** Prepared medium is trace to slightly hazy and pale to light yellow.

**Expected Cultural Response:** Cultural response on YM Agar at 25 - 30°C after 2 - 7 days incubation.

Microorganism	Response	Reactions
<i>Aspergillus niger</i> ATCC® 16404	growth	white cottony to black powdery
<i>Candida albicans</i> ATCC® 10231	growth	off-white to beige pasty
<i>Lactobacillus fermentum</i> ATCC® 9338	growth	---
<i>Microsporium canis</i> ATCC® 36299	growth	white cottony-powdery
<i>Penicillium roquefortii</i> ATCC® 10110	growth	grey powdery
<i>Saccharomyces cerevisiae</i> ATCC® 9763	growth	off-white to beige, pasty

The organisms listed are the minimum that should be used for quality control testing.

### **Test Procedure**

1. Inoculate YM Agar with the appropriate sample for the presence of yeasts, molds, or aciduric microorganisms.
2. Incubate at 30 ± 2°C for 2 to 7 days.

### **Results**

Examine plates for growth. Record YM Agar results as colony forming units (CFU) per volume of sample.

### **Storage**

Store sealed bottle containing the dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

### **Expiration**

Refer to expiration date stamped on container. The dehydrated medium should be discarded if not free flowing, or if appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### **Limitations of the Procedure**

1. Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.
2. Acidified YM Agar should not be reheated.

### **Packaging**

<b>YM Agar</b>	<b>Code No.</b>	<b>7525A</b>	<b>500 g</b>
		<b>7525B</b>	<b>2 kg</b>
		<b>7525C</b>	<b>10 kg</b>

### **References**

1. 1951. U. S. Dept. Agricult. Tech. Bull. No. 1029.
2. 1939. J. Tropical Med. Hyg. 42:176.
3. Jong, S. C., and M. J. Edwards. 1991. American Type Culture Collection Catalog of filamentous fungi. 18<sup>th</sup> ed. American Type Culture Collection, Rockville, MD.

### **Technical Information**

Contact Acumedia Manufacturers, Inc. for Technical Service or questions involving dehydrated culture media preparation or performance at (517)372-9200 or fax us at (517)372-2006.